

The central density of neutron stars in close binaries

Alan G. Wiseman

*Theoretical Astrophysics, 130-33, California Institute of Technology
Pasadena, CA 91125*

E-mail: agw@tapir.caltech.edu

(February 7, 2008)

Recent numerical simulations of coalescing binary neutron stars conducted by Wilson, Mathews and Marronetti (WMM) show a rising central energy density of the stars as the orbital separation shrinks, *i.e.* the stars are individually crushed as they near coalescence. They claim this “star-crushing” effect is partially due to a non-linear, first post-Newtonian order enhancement of the self-gravity of each star caused by the presence of the other star. We present a concrete calculation which shows, within general relativity, *first post-Newtonian order interactions with the other star leave the central energy density unchanged as the orbital radius shrinks*. The results presented here are in sharp disagreement with the WMM claim. However, alternative gravitational theories, such as Brans-Dicke theory, can exhibit a small crushing effect in the binary constituents as they near coalescence. We show that the absence of the star-crushing effect at first post-Newtonian order is related to adherence to the strong equivalence principle.

PACS Numbers: 97.80.Fk, 04.25.Dm, 04.40.Dg, 97.60.Jd

Binary neutron stars spiraling toward their final coalescence under the dissipative influence of gravitational radiation reaction forces are the primary sources gravitational waves which might be detected by interferometric gravitational wave detectors such as LIGO and VIRGO [1,2]. Extracting the gravitational waves from the detector noise and using the information encoded in the signals requires a precise knowledge of the expected waveforms [2]. The primary method for studying the behavior of these highly relativistic binaries has been the “post-Newtonian” approximation carried to high relativistic order [3–6]. Recently, Wilson, Mathews and Marronetti (WMM) [7] have used numerical hydrodynamic simulations as an alternative method for studying the late stages of binary neutron-star inspiral. These simulations, based on the simplifying assumption that space-time is spatially conformally-flat (SCF) [8], obtain a sequence of equilibrium configurations of the neutron stars at closer and closer orbital separation. Their numerical results show that the neutron stars are individually compressed and the central densities begin to rise as the orbital separation is decreased. [See Mathews and Wilson [7] Figure 2.] Such a crushing-effect has never been reported in post-Newtonian calculations. This discrepancy forces a re-examination of the validity of both the post-Newtonian calculations and the numerical simulations. (See Eardley [9] for a discussion.)

It is tempting to simply dismiss this unusual star-crushing effect as an artifact of the SCF constraint used in the WMM numerical simulations. However, WMM further claim that the star-crushing effect is partially due to a non-linear first post-Newtonian order (1PN) enhancement of the gravitational potential. (See Mathews and Wilson [7] Appendix A: “... the [first] post Newtonian expansion does indeed exhibit an enhancement to the gravitational potential ...”) However, the 1PN metric is naturally spatially conformally flat; therefore such an effect would be independent of the SCF assumption. [See Eq.(1).] It is also tempting to dismiss the WMM claim on the grounds that Newtonian tidal-effects will stretch the individual stars, and thereby reduce the central densities of stars [10]. However, WMM have made it clear that their proposed crushing mechanism will swamp – by a factor of 10^4 – the rather small reduction in central density due to tidal distortion. [See Mathews and Wilson [7] Eq. (20).]

There is actually a substantial body of literature on post-Newtonian effects for extended bodies in binaries that seemingly would have discovered a 1PN star-crushing effect if it were real. (See *e.g.* [11,12].) There are also published arguments that show that enhancements of the local gravity due the presence of other gravitating bodies involve a 1PN violation of the strong equivalence principle. (See Will [3] Chapter 6.3.) However, WMM specifically claim a 1PN origin of the star-crushing effect; therefore we offer a specific rebuttal to the claim.

In this paper we present a brutally straightforward, first post-Newtonian order calculation of the central energy density of a compact star in a close binary. We solve the 1PN hydrodynamic equation (including non-linear gravitational interactions with the other star) for the density profile of the star. At intermediate steps in the calculation, relativistic corrections appear which give the illusion that each star is being compressed by the presence of the other star. In the end, for a first post-Newtonian expansion of general relativity, all the relativistic corrections involving the other star cancel, and the true physical quantities (the central energy density, and the proper radius) are unaffected by the presence of the other star. Thus WMM’s claim of a 1PN origin of the star-crushing effect is false. The confusion can be summed up as follows: the potential [in this case ($g_{00} + 1$)] is made “deeper” by non-linear relativistic factors involving the other star; thus, when solving for the density profile, it appears that the star is being com-

pressed by the presence of the other star. However, the relativistic factors also re-enter the calculation as 1PN corrections to the spatial part of the metric. The spatial metric components alter the spatial line element in the calculation of the proper radius of the star, and they alter the proper volume element in the calculation of the conserved baryon number. In general relativity, at first post-Newtonian order, these effects exactly cancel, and the physical quantities are left unaffected by the other star.

We consider two perfect fluid blobs (stars) with baryon masses M_{bA} and M_{bB} . For simplicity we assume that the stars are in a nearly circular orbit with coordinate radius R_o . We boost to a comoving, post-Newtonian coordinate system where the fluid in star-A is *momentarily* at rest, and the origin instantaneously coincides with the center of star-A. [Star-A is still being accelerated toward star B.] In this moving frame the parameterized, post-Newtonian (PPN) metric coefficients are [3,13]

$$g_{00} = -1 + 2U(\mathbf{x}, t) - 2\beta U(\mathbf{x}, t)^2 + 2(\gamma + 1)\Phi_1(\mathbf{x}, t) + 2(3\gamma - 2\beta + 1)\Phi_2(\mathbf{x}, t) + 2\Phi_3(\mathbf{x}, t) + 6\gamma\Phi_4(\mathbf{x}, t) \quad (1a)$$

$$g_{ij} = [1 + 2\gamma U(\mathbf{x}, t)]\delta_{ij} \quad (1b)$$

$$g_{0j} = -(1/2)(4\gamma + 3)V_j(\mathbf{x}, t) - (1/2)W_j(\mathbf{x}, t), \quad (1c)$$

where β and γ are the PPN coefficients which characterize alternative theories of gravity. Solar system tests have constrained their values to be very close to unity. In general relativity $\beta \equiv \gamma \equiv 1$. Also notice the metric is spatially conformally flat (SCF) [8]. The sources for the potentials (the baryon density ρ , pressure P , and internal thermal energy $\rho\Pi$) are measured in the comoving frame of the fluid.

We now define and evaluate the potentials appearing in Eq.(1). (Details can be found in [3].) The Newtonian potential is

$$U(\mathbf{x}, t) \equiv \int \frac{\rho(\mathbf{x}', t)d^3\mathbf{x}'}{|\mathbf{x} - \mathbf{x}'|} = \frac{M_{bB}}{R_o} + \frac{M_{bB}\mathbf{R}_o \cdot \mathbf{x}}{R_o^3} + 4\pi \left[\frac{1}{r} \int_0^r \rho_A(r') r'^2 dr' + \int_r^L \rho_A(r') r' dr' \right]. \quad (2)$$

where L denotes the – as yet unknown – coordinate radius of the star, $\mathbf{R}_o$ is the vector separation of the body centers, $R_o = |\mathbf{R}_o|$ is the constant coordinate orbital separation, $\mathbf{x}$ is the field point, $r = |\mathbf{x}|$ is the radial coordinate measured from the center of star-A. The second expression in Eq.(2) is valid only within the quasi-stationary star-A. We have also neglected tidal terms. The $(\mathbf{R}_o \cdot \mathbf{x})$ -term gives rise to the bulk acceleration of star-A toward star-B. Over the extent of star-A, it has zero average radial (r) component; therefore it cannot contribute to the compression. We will neglect this term and other terms which clearly cannot contribute to the radial compression. The Newtonian potential can now be written

$$U(r < L) = \frac{M_{bB}}{R_o} + U_A(r), \quad (3)$$

where $U_A(r)$ is defined as the “ 4π ”-term in Eq. (2). For future reference, we compute the radial derivative

$$U_{,r} = -\frac{4\pi}{r^2} \int_0^r \rho(r') r'^2 dr' \equiv -\frac{M_{bA}(r)}{r^2}. \quad (4)$$

The potential Φ_1 is the post-Newtonian contribution from the kinetic energy. As we have boosted to a frame where star-A is at rest, there is no bulk motion of the fluid in star-A to contribute to this term. We also neglect internal fluid motion in star-A, as this actually decreases the central density [10]. Thus we have

$$\Phi_1(\mathbf{x}, t) = \int_B \frac{v^2 \rho(\mathbf{x}', t) d^3\mathbf{x}'}{|\mathbf{x} - \mathbf{x}'|} = v_B^2 \frac{M_{bB}}{R_o}, \quad (5)$$

where v_B is the velocity of star-B measured in the frame where star-A is momentarily at rest. The potential Φ_2 is the post-Newtonian contribution from the gravitational energy

$$\Phi_2(\mathbf{x}, t) \equiv \int \frac{U(\mathbf{x}', t)\rho(\mathbf{x}', t)d^3\mathbf{x}'}{|\mathbf{x} - \mathbf{x}'|}. \quad (6)$$

Proceeding as with previous terms we have

$$\Phi_{2,r} = -\frac{M_{bB}}{R_o} \frac{M_A(r)}{r^2} - \frac{4\pi}{r^2} \int_0^r \rho_A(r') U_A(r') r'^2 dr'. \quad (7)$$

The origin of the WMM star-crushing claim is that the first term in Eq.(7) will conspire with the Newtonian gravitational pull in Eq.(4) to enhance the compression of star-A. Noting that the WMM-factor $(W^2 - 1) \sim M_{bB}/R_o$, we see that radial enhancement to the Newtonian compressional force appearing in Eq.(7) is the force related to the potential energy shown in Mathews and Wilson [7] Eq.(19) [$\Delta E_{GR} \approx 2(W^2 - 1)GM^2/L$]. Thus, we are unambiguously addressing the term that WMM claim is the origin of the crushing-effect.

The potentials Φ_3 and Φ_4 are similar to Φ_1 with the kinetic energy replaced by $\rho\Pi$ and P respectively. Finally, the potential can be evaluated using formulae in [3] The results: $V_j = M_{bB}v_B^j/R_o$, and $W_j = 0$ for circular orbits.

We now look at the spatial components of the Euler equation in our instantaneously comoving-moving frame of star-A. In this frame, the fluid is momentarily stationary ($u^k = P_{,t} = 0$ for star-A), and we can write

$$\frac{dP}{dr} = -(\rho + \rho\Pi + P) \left[\frac{d}{dr} \log \sqrt{-g_{00}} \right]. \quad (8)$$

We have neglected a number or second post-Newtonian terms, as well as terms which do not contribute to the radial compression.

Expanding the logarithm, substituting the potentials, and discarding second post-Newtonian terms, we have

$$\frac{dP}{dr} - \rho U_{,r} = \epsilon \rho \left[2(1 - \beta)UU_{,r} + (3\gamma - 2\beta + 1)\Phi_{2,r} + \Phi_{3,r} + 3\gamma\Phi_{4,r} + [\Pi + (P/\rho)]U_{,r} \right]. \quad (9)$$

On the left we have the familiar pressure gradient balancing the Newtonian gravitational attraction. On the right we have all the first post-Newtonian corrections. We have inserted a dummy factor of $\epsilon (= 1)$ to keep track of post-Newtonian terms. Multiplying by r^2/ρ , differentiating, and using the expressions for the radial derivatives of the potentials, we have

$$\begin{aligned} & \frac{1 - \epsilon(3\gamma - 4\beta + 3)(M_{bB}/R_o)}{4\pi} \frac{1}{r^2} \frac{d}{dr} \left[\frac{r^2}{\rho} \frac{dP_A}{dr} \right] + \rho_A \\ &= -\epsilon \left\{ \frac{\beta - 1}{2\pi r^4} M_A(r)^2 + \rho_A [(3\gamma - 4\beta + 3)U_A + \Pi_A] \right. \\ & \quad \left. + (3\gamma + 1)P_A + \frac{d}{dr} \left(\Pi_A + \frac{P_A}{\rho_A} \right) \frac{M_A(r)}{4\pi r^2} \right\}. \quad (10) \end{aligned}$$

We have discarded terms of second post-Newtonian order, *i.e.* $O(\epsilon^2)$. On the left of Eq.(10) we have Newtonian terms, as well as the post-Newtonian terms related to the presence of star-B. On the right we have only post-Newtonian self-interaction terms. The terms on the right will affect the matter configuration of star-A, but they would do so whether or not star-B is nearby. Our focus is on deriving the 1PN effect star-B has on the matter configuration in star-A. This can be done by setting the right hand side of Eq. (10) to zero and solving the homogeneous equation. In a full solution to Eq.(10) any non-linear interplay between the post-Newtonian corrections on the left and right sides of Eq.(10) would be $O[\epsilon^2]$, and could be neglected. Throughout the remainder of this calculation, we ruthlessly discard post-Newtonian self interaction terms. (A 1PN solution of Eq.(10) – including the self interactions on the right side – will be presented elsewhere [14].)

Substituting the polytropic equation of state

$$P = K\rho^{1+1/n}, \quad (11)$$

into the homogeneous equation, and defining the quantities

$$\rho = \rho_c \theta^n \quad (12a)$$

$$a = \sqrt{\frac{(n+1)K}{4\pi}} \quad (12b)$$

$$\lambda = a\rho_c^{\frac{1-n}{2n}} \left[1 - \frac{\epsilon}{2} \frac{M_{bB}}{R_o} (3\gamma - 4\beta + 3) \right] \quad (12c)$$

$$\xi = r/\lambda. \quad (12d)$$

we obtain the dimensionless Lane-Emden [15] equation

$$\frac{1}{\xi^2} \frac{d}{d\xi} \left[\xi^2 \frac{d\theta}{d\xi} \right] + \theta^n = 0. \quad (13)$$

[We restrict our attention to non-singular ($n < 3$) solutions [15].] The solutions $\theta(\xi; n)$ of this non-linear differential equation must satisfy the boundary conditions

$$\theta(0; n) = 1, \quad \theta'(0; n) = 0. \quad (14)$$

The first condition defines ρ_c as the central baryon density. The second insures there is no pressure gradient at the origin. The density profile of star-A can then be written

$$\rho_A(r) = \rho_c \theta(r/\lambda; n)^n. \quad (15)$$

Notice that $\rho_A(r)$ has an apparent 1PN dependence on the potential produced by the other star (M_{bB}/R_o) through Eq.(12c). The coordinate radius L of the stellar surface is the value of r where the pressure (and thereby the density) vanishes; it is given by

$$L = \lambda \xi_{1n}, \quad (16)$$

where ξ_{1n} is the smallest positive root satisfying $\theta(\xi_{1n}; n) = 0$.

The total baryon mass can be obtained by integrating

$$M_{bA} = \int_A \rho d(\text{proper volume}) = \int_A \rho u^0 \sqrt{-g} d^3x \quad (17a)$$

$$= \int_A \rho [1 + 3\epsilon\gamma U] d^3x \quad (17b)$$

$$= 4\pi a^3 f(n) \rho_c^{\frac{3-n}{2n}} \left[1 + \frac{3}{2} \frac{M_{bB}}{R_o} (4\beta - \gamma - 3) \right] \quad (17c)$$

$$= M_{bA} [K, n, \rho_c, (4\beta - \gamma - 3)(M_{bB}/R_o)] \quad (17d)$$

$$= 4\pi a^3 f(n) \rho_c^{\frac{3-n}{2n}} \quad (\text{in gen. rel. } \beta = \gamma = 1), \quad (17e)$$

$$= M_{bA} [K, n, \rho_c] \quad (\text{in general relativity}), \quad (17f)$$

where $f(n) = \xi_{1n}^2 |\theta'(\xi_{1n}, n)|$ is solely a function of the polytropic index n . We have discarded self-interaction terms and 2PN terms. (We have also set the dummy $\epsilon = 1$.) *Notice, – and this is the point of the paper – in general relativity at 1PN order for a star with fixed baryon mass (M_{bA}) and a fixed equation of state (K and n), there will be no change in the central density as the binary orbital separation changes.* The functional dependence on M_{bB}/R_o cancels out of Eq. (17) for general relativity (or any gravitational theory with $4\beta - \gamma - 3 = 0$). The inclusion of the self interactions that we have been discarding would complicate the functional form shown Eq. (17e), but it would not affect the the functional dependence depicted in Eq. (17f); therefore our *no-crushing* conclusion would hold even if we were to include the self interactions on the right of Eq (10).

The proper radius, Δ_A , of the star can be obtained by integrating

$$\Delta_A = \int_0^L \sqrt{g_{rr}} dr \quad (18a)$$

$$= L_N \left[1 + \frac{n}{3-n} \frac{M_{bB}}{R_o} (4\beta - \gamma - 3) \right] \quad (18b)$$

$$= L_N \quad (\text{in general relativity } \beta \equiv \gamma \equiv 1). \quad (18c)$$

where

$$L_N = a \xi_{1n} \left(\frac{M_{bA}}{4\pi a^3 f(n)} \right)^{\frac{1-n}{3-n}} \quad (19)$$

is the radius of a the isolated Newtonian star with baryon mass M_{bA} . *Notice, in general relativity at 1PN order, the proper radius of the star in a close binary is unaffected by the presence of the other star.*

In the Brans-Dicke theory of gravity $\beta \equiv 1$ and $|\gamma-1| \lesssim 0.001$ [3]; therefore the compression effect illustrated in Eq. (17) and (18) will be small. However, even with these constrained values of β and γ , this effect is potentially larger than those due to tidal distortion. [See Mathews and Wilson [7] Eq.(20).]

The 1PN crushing effect we have been discussing can be interpreted as a change in the local value of Newton's gravitational constant due to the presence of the other star in the binary. That is, Eq. (17c) and Eq. (18b) can be obtained from the standard Newtonian formula [15] by the replacing $G = 1$ with

$$G_{local} = 1 - (4\beta - \gamma - 3)(M_{bB}/R_o) . \quad (20)$$

Such effects arise in gravitational theories that violate the strong equivalence principle. [See reference [3] Eq. (6.75).]

Notice, we have only shown that there is no crushing effect at 1PN order in general relativity. As the conventional (perfect fluid) post-Newtonian expansion of the metric is naturally spatially conformally flat at 1PN order, we cannot conclude with certainty that the crushing effect that WMM see is an artifact of the SCF constraint used in the numerical simulations. Indeed, it is possible that the crushing effect is real, and WMM's only sin was to incorrectly assign a 1PN origin to the effect. The question remains whether the SCF boundary condition is somehow responsible for (or amplifies) the crushing effect in the simulations [16].

The post-Newtonian expansion of the metric is not spatially conformally flat at second post-Newtonian order [17,18]; therefore a thorough 2PN investigation may clarify the difference between the conventional post-Newtonian analysis of the inspiral problem and the SCF numerical analysis [14]. WMM claim that the SCF constraint is justified, because it removes only the radiation from the solution. This is not true. Radiation only enters the metric at 2.5PN order; thus altering the metric at 2PN order by imposing the SCF constraint will change the longitudinal fields that govern the periodic orbital motion and the individual body configurations. Although it is true that the WMM equilibrium solutions are exact solutions (up to numerical accuracy) of the Einstein field equations, they are solutions which differ from the actual inspiral metric at the 2PN level. In this formal sense, the SCF simulations are only as accurate as a 1PN simulation.

It is a pleasure to thank Patrick Brady, Scott Hughes, Lee Lindblom, Katrin Shenk and Clifford Will for their helpful insights. This work was supported by NSF grant PHY-9424337 and AST-9417371.

-
- [1] A. Abramovici, et al., Science, **256**, 325 (1992).
 - [2] K. S. Thorne, in "Proceedings of Snowmass 94 Summer Study on Particle and Nuclear Astrophys. and Cosmology", eds. E. W. Kolb and R. Peccei (World Scientific, Singapore, 1995).
 - [3] C.M. Will, *Theory and Experiment in Gravitational Physics* (Cambridge University Press, Cambridge, England, 1981). See particularly Table 4.1 and 6.1. We use the notation and conventions of this reference. In particular we use unit in which $G = c = 1$, and spatial indices are denoted $i, j, k, \dots$
 - [4] L. Blanchet, T. Damour, B. R. Iyer, C. M. Will and A. G. Wiseman, Phys. Rev. Lett., **74**, 3515 (1995).
 - [5] C. M. Will, and A. G. Wiseman, Phys. Rev. D **54** 4813 (1996).
 - [6] Loosely speaking, the post-Newtonian expansion is a Taylor series approximation to general relativity, or any alternative theory of gravity. At first post-Newtonian order (1PN) we keep relativistic corrections of $O[(v/c)^2] \approx O[(GM/rc^2)]$ beyond the leading order Newtonian terms. We discard 2PN terms (*e.g.* $O[(GM/rc^2)^2]$) throughout this paper.
 - [7] J. R. Wilson and G. J. Mathews, Phys. Rev. Lett. **75**, 4161 (1995); J. R. Wilson, G. J. Mathews and P. Maronetti, Phys. Rev. D. **54**, 1317; G.J. Mathews and J.R. Wilson astro-ph/9701142. These three papers are referred to as WMM.
 - [8] For the uninitiated, spatial conformal flatness means the spatial part of the metric can be written as $g_{ij} = (\text{conformal factor}) \times \delta_{ij}$.
 - [9] D. Eardley in *Matters of Gravity* gr-qc/9702010
 - [10] D. Lai, Phys. Rev. Lett. **76**, 4878 (1996).
 - [11] A. G. Wiseman, Phys. Rev. D **46** 1517 (1992).
 - [12] L. Blanchet and G. Schäfer, Mon. Not. R. Astron. Soc. **239**, 845 (1989).
 - [13] Virtually all Newtonian gravitational effects can be obtained from the metric coefficients $g_{00} = -1 + 2U$, $g_{0i} = 0$, and $g_{ij} = \delta_{ij}$. We are including "post"-Newtonian (1PN) corrections to all the metric coefficients.
 - [14] A.G. Wiseman, In preparation.
 - [15] S. Chandrasekhar, *An Introduction to the Study of Stellar Structure* (University of Chicago Press, Chicago, IL, 1939).
 - [16] Keep in mind that the imposition of the SCF constraint is not an "alternative theory of gravity" (in the sense that Brans-Dicke is an alternative theory). Rather, the SCF constraint is really a stringent boundary condition on the solutions to the actual general relativistic field equations.
 - [17] C.M. Will, private communication.
 - [18] R.Reith and G. Schäfer, preprint gr-qc/9603043.